

Molecular comparison of topotypic specimens confirms *Anopheles (Nyssorhynchus) dunhami* Causey (Diptera: Culicidae) in the Colombian Amazon

Freddy Ruiz^{1,2,3/+}, Yvonne-Marie Linton², David J Ponsonby³, Jan E Conn⁴,
Manuela Herrera⁵, Martha L Quiñones⁵, Iván D Vélez⁶, Richard C Wilkerson¹

¹Division of Entomology, Walter Reed Army Institute of Research, 503 Robert Grant Av., 20910 Maryland, USA ²Mosquitoes Programme, Department of Entomology, Natural History Museum, London, UK ³Department of Geographical and Life Sciences, Canterbury Christ Church University, Kent, UK ⁴Griffin Laboratory, Wadsworth Center, New York State Department of Health, Albany, NY, USA

⁵Facultad de Medicina, Universidad Nacional de Colombia, Bogotá, Colombia ⁶Programa de Estudio y Control de Enfermedades Tropicales, Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia

The presence of Anopheles (Nyssorhynchus) dunhami Causey in Colombia (Department of Amazonas) is confirmed for the first time through direct comparison of mtDNA cytochrome c oxidase I (COI) barcodes and nuclear rDNA second internal transcribed spacer (ITS2) sequences with topotypic specimens of An. dunhami from Tefé, Brazil. An. dunhami was identified through retrospective correlation of DNA sequences following misidentification as Anopheles nuneztovari s.l. using available morphological keys for Colombian mosquitoes. That An. dunhami occurs in Colombia and also possibly throughout the Amazon Basin, is of importance to vector control programs, as this non-vector species is morphologically similar to known malaria vectors including An. nuneztovari, Anopheles oswaldoi and Anopheles trinkae. Species identification of An. dunhami and differentiation from these closely related species are highly robust using either DNA ITS2 sequences or COI DNA barcode. DNA methods are advocated for future differentiation of these often sympatric taxa in South America.

Key words: *Anopheles dunhami* - COI barcodes - ITS2 - topotypic - Brazil - Colombia

Anopheles (Nyssorhynchus) dunhami Causey was first described from Tefé, Amazonas (AM), Brazil (Causey 1945), but was placed in synonymy with the malaria vector *Anopheles nuneztovari* Gabaldón (Lane 1953), where it remained for some decades until taxonomic revisions resulted in its re-elevation (Peyton 1993). In the same study, Peyton also reduced the widespread east-Andean taxon *Anopheles trinkae* Faran to a junior synonym of *An. dunhami*, based on similarities in male genitalia (Peyton 1993). Later integrated studies, combining egg morphology, cytogenetics and rDNA sequences of the second internal transcribed spacer (ITS2), confirmed that *An. trinkae* and *An. dunhami* were indeed separate taxa and *An. trinkae* was reinstated to species status (Lounibos et al. 1998). Recently, Calado et al. (2008) reaffirmed the differences between *An. dunhami* and *An. nuneztovari* using the morphology of male genitalia and DNA sequences of the ITS2 and a portion of the mitochondrial cytochrome c oxidase I (COI). Overlapping morphological characters between proven malaria vec-

tors *An. nuneztovari*, *An. trinkae* and *Anopheles oswaldoi* (Hayes et al. 1987, Olano et al. 2001, Quiñones et al. 2006, Gutiérrez et al. 2009) and isomorphic non-vector species such as *An. dunhami* continue to hamper correct species identification in South America and can impact the effectiveness of vector control programs.

As a result of its historically confused taxonomic status, few verified distribution records exist for *An. dunhami*. Records to date are limited to AM: Tefé (type locality) (Causey 1945, Peyton 1993), Tabatinga (Lounibos et al. 1998) and Parintins (Calado et al. 2008). Although Tabatinga is adjacent to the Colombian border, *An. dunhami* has never been reported from Colombia and the species is not included in the available morphological keys to Colombian taxa (Suárez et al. 1988, González & Carrejo 2007). Herein, we molecularly characterize topotypic *An. dunhami* using COI barcodes [after Hebert et al. (2003)] and ITS2 rDNA sequences and, through molecular correlation of sequence data, confirm the presence of *An. dunhami* in the Colombian Amazon for the first time.

MATERIALS AND METHODS

Mosquito specimens - A total of 4,893 mosquitoes were collected in four departments in Colombia (Antioquia, Amazonas, Caquetá and Norte de Santander) in 2006 as part of the PhD study of Ruiz (2010). Of 574 mosquitoes collected in Amazonas, where *An. nuneztovari* s.l. is unreported, two specimens were keyed out as *An. nuneztovari* s.l. using the morphological keys of Suárez et al. (1988) and Faran (1980). DNA was extracted

Financial support: UNICEF/UNDP/WorldBank/WHO/TDR (A50252 to YML), Canterbury Christ Church University, UK (PhD studentship for FR to DJP and YML), NIH, USA (2R01AI054139 to JEC), Smithsonian Institution, Walter Reed Army Institute of Research + Corresponding author: ruizj@si.edu

Received 29 April 2010

Accepted 22 July 2010

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE APR 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Molecular comparison of topotypic specimens confirms Anopheles (Nyssorhynchus) dunhami Causey (Diptera: Culicidae) in the Colombian Amazon				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Walter Reed Biosystematics Unit, 503 Robert Grant Avenue, Silver Spring, MD, 20910				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

TABLE

Field codes, collection localities and co-ordinates of specimens of *Anopheles dunhami* and *Anopheles nuneztovari* C used in this study, alongside GenBank accessions for second internal transcribed spacer (ITS2) and cytochrome c oxidase I (COI) sequences where appropriate outgroup taxa were chosen based on their close morphological similarity to *An. dunhami*

Species	Country, locality (co-ordinates)	Field code	GenBank			
			ITS2	COI		
<i>An. dunhami</i>	Brazil, Amazonas, Tefé (-03.3207, -64.7235)	SA26	HQ020394	-		
		SA44	HQ020395	HQ315878		
		SA79	HQ020396	HQ315879		
		SA85	HQ020397	HQ315880		
		SA98	HQ020398	HQ315881		
	Colombia, Amazonas, Leticia (-04.1158, -69.9522)	C08921	HQ020382	-		
		C09041	HQ020383	HQ315867		
	<i>An. nuneztovari</i> C	Colombia, Antioquia, Zaragoza (07.4847, -74.8679)	Za-106	HQ020399	-	
			Za-110	HQ020400	-	
			Za-130	HQ020401	-	
Za-131			HQ020402	-		
Za-158			HQ020403	-		
Za-207			HQ020404	-		
Za-208			HQ020405	-		
Colombia, Norte de Santander, Tibú (08.6403, -72.7372)			NS-17	HQ020388	HQ315872	
		NS-62	HQ020391	HQ315875		
		NS-71	HQ020392	HQ315876		
		NS-72	HQ020393	HQ315877		
		NS-130	HQ020384	HQ315868		
		NS-135	HQ020385	HQ315869		
		NS-139	HQ020386	HQ315870		
		NS-144	HQ020387	HQ315871		
		NS-243	HQ020389	HQ315873		
		NS-319	HQ020390	HQ315874		
		Outgroups				
		<i>Anopheles oswaldoi</i> B	Colombia, Antioquia, Nechí (08.1100, -74.7670)	CO9000	-	HQ315866
<i>Anopheles rangeli</i>		Brazil, Mato Grosso (-10.2256, -54.9861)	BR704-40	-	HQ315865	
<i>Anopheles trinkae</i>	Ecuador, Napo, Sardina Yacu (-00.1666, -77.0833)	SY205	-	HQ315882		
		SY227	-	HQ315883		

from these two specimens and a further 17 individuals of *An. nuneztovari* from the departments of Zaragoza (n = 7) and Norte de Santander (n = 10) (Table). Given the morphological similarities between *An. nuneztovari* and

An. dunhami and the reported presence of the latter species in the Brazilian Amazon, DNA was also extracted from abdomens of five archive specimens of *An. dunhami* collected at the type locality (Tefé, Brazil). Voucher

specimens are stored in the mosquito collections of the Smithsonian Institution of National Museum of Natural History (NMNH), Washington DC, USA (Table).

Molecular methods - DNA template was acquired using the commercially available DNeasy Blood & Tissue Kit and BioSprint 96 DNA (both QIAgen®, Maryland, USA). Amplification of rDNA ITS2 polymerase chain reaction (PCR) was carried out using the primers of Collins and Paskewitz (1996) and the protocol described by Linton et al. (2001).

Amplification of the 710 bp fragment of the barcoding region of the mitochondrial COI gene was achieved using the primer pair LCO 1490 and HCO 2198 developed by Folmer et al. (1994), using a PCR protocol optimized by the Laboratories of Analytical Biology, Museum Support Center, Smithsonian Institution/NMNH. Each PCR contained 1 x NH₄ buffer, 0.5 mM each dNTP, 2 mM of MgCl₂, 0.2 U of Taq polymerase (BioLine, Taunton, MA, USA), 0.3 µM each primer and 1 µL of DNA template and was made up to a total volume of 10 µL using ddH₂O. The PCR thermocycler parameters included: a single cycle at 95°C for 5 min, followed by 34 cycles of 95°C for 30 s, 48°C for 30 min and 72°C for 45 s, respectively, terminating with a 72°C for 5 min extension step and a 10°C hold. PCR products (both ITS2 and COI) were visualized on 1% agarose gels, containing 0.5 mg/mL of ethidium bromide and purified using ExoSAP-IT® (USB Corporation, Cleveland, Ohio, USA).

Sequencing reactions were carried out in both directions using the Big Dye Terminator Kit® (PE Applied BioSystems, Warrington, England) on an ABI 3730 automated sequencer (PE Applied BioSystems). Sequences were edited using Sequencer™ v.4.8 (Genes Codes Corporation, Ann Arbor, MI) and either aligned automatically in ClustalX (Thompson et al. 1997) or manually using MacClade v.4.06 (Maddison & Maddison 2003). Sequence similarities were compared with those available in GenBank using Basic Local Alignment Search Tool (blast.ncbi.nlm.nih.gov/Blast.cgi). Sequence divergence was calculated in MEGA version 4 (Kumar et al. 2008) using the Kimura 2-parameter distance model (K2P) (Kimura 1980). A bootstrap neighbor joining (NJ) tree (Saitou & Nei 1987) was generated in PAUP version 4.0 (Swofford 2003), using 1,000 replicates of the K2P distance matrices.

COI sequences generated formed part of the activities of the Mosquito Barcoding Initiative (project leader YML) and both full specimen and COI sequence data, including electropherograms, are freely available through the Barcode of Life database website at boldsystems.org. GenBank accession for sequences generated in this study are listed in Table.

RESULTS

ITS2 sequences (n = 5) and COI DNA barcodes (n = 4) were generated from five topotypic archive specimens of *An. dunhami* from Tefé. ITS2 sequences from 19 Colombian *Anopheles*, morphologically identified as *An. nuneztovari* using the keys of Faran (1980) and Suárez et al. (1988) revealed two distinct ITS2 sequenc-

es, differing at 17 bases [1 transition (G/A), 1 transversion (C/G) and 15 indels] (Fig. 1).

The most frequent ITS2 sequence was shared by seven specimens from Zaragosa (Antioquia) and 10 specimens from Tibú (Norte de Santander), morphologically identified as *An. nuneztovari* s.l., whereas a different sequence was shared by two specimens collected in Kilómetro 12, Leticia (Amazonas, Colombia). The most frequent sequence was identical to previously published sequences of *An. nuneztovari* C from Colombia (Sierra et al. 2004) (AY028081-124, AY028128), Brazil (Marrelli et al. 2005) (AF461749) and Venezuela (Fritz et al. 1994) (L22462).

The two specimens from Kilómetro 12, Leticia showed 100% identity with topotypic *An. dunhami* sequences and three published *An. dunhami* sequences from Parintins (EU848337, EU848338 and EU848344) (Calado et al. 2008). However, sequence differences were noted between the topotypic and Colombian specimens and the remaining six available *An. dunhami* sequences (generated from clones) reported from Parintins. Topotypic and Colombian *An. dunhami* shared 99.7% identity with EU848340 and EU848341 (1/360 variant bases), 99.4% identity with EU848339, EU848343, EU84845, same haplotype (2/360 variant bases) and 98.9% with EU848342 (4/360 variant bases) (Fig. 1).

MtDNA COI DNA barcode sequences (658 bp without primers) were generated for specimens previously identified by ITS2 sequence data as follows: topotypic *An. dunhami* from Brazil (n = 4) and Colombia (Kilómetro 12, Leticia) (n = 1) and *An. nuneztovari* C from Norte de Santander (n = 10). COI barcode data were also generated for *An. trinkae* (n = 2), *An. oswaldoi* B (n = 1) and *An. rangeli* (n = 1), as these species can be morphologically mistaken for *An. dunhami* or *An. nuneztovari* s.l. in the female adult stage (Faran 1980).

The bootstrap NJ tree (Fig. 2), based on 1,000 replicates of the K2P COI distance matrices (Kimura 1980), reaffirmed the specific identity of topotypic and Colombian *An. dunhami* and its genetic differentiation from other closely related taxa including *An. nuneztovari* C, *An. trinkae*, *An. oswaldoi* B and *An. rangeli* obtained by ITS2. Mean intra-specific K2P

		122222222233333333
		292247888889011225555
		492339012340845453456
<i>An. dunhami</i> (n=5), Bra., Tefé		t---aagagagaaga-----
<i>An. dunhami</i> (n=2), Col., Amazonas	
<i>An. dunhami</i> (EU848337-38,44), Bra., Par.	
<i>An. dunhami</i> (EU848339,43,45), Bra., Par.	
<i>An. dunhami</i> (EU848340), Bra., Par.		---g-----
<i>An. dunhami</i> (EU848341), Bra., Par.	g-----
<i>An. dunhami</i> (EU848342), Bra., Par.		c-----c-----
<i>An. nuneztovari</i> (n=17), Col., N.S., Zar.		.ggg.-----gc.tgagag

Fig. 1: alignment showing differing bases of second internal transcribed spacer sequences of Colombian *Anopheles dunhami* (546 bp; n = 2) and *Anopheles nuneztovari* C (549 bp; n = 17) generated in this study in comparison to topotypic *An. dunhami* from Tefé, Brazil (546 bp; n = 5) and other available sequences available in GenBank (360 bp; EU848337-42) (Calado et al. 2008). Bra: Brazil; Col: Colombia; Par: Parintins; N.S.: Norte de Santander; Zar: Zaragoza.

divergence values were 0.007, 0.004 and 0.010 in *An. dunhami* (n = 5), *An. nuneztovari* C (n = 10) and *An. trinkae* (n = 2), respectively. The mean K2P value between *An. dunhami* and *An. nuneztovari* was 0.025, *An. dunhami* and *An. trinkae* was 0.093, *An. nuneztovari* and *An. trinkae* was 0.085.

DISCUSSION

This study provides DNA sequence data for topotypic specimens of *An. dunhami* for the first time, thus establishing the genetic identity of the species. Correlation of the ITS2 rDNA and COI barcodes of Colombian specimens from the Colombian Amazon with topotypic *An. dunhami* confirms the first record of *An. dunhami* outside Brazil and comprises a new record for Colombia. That *An. dunhami* can be morphologically misidentified with vector species, including *An. nuneztovari* s.l. (as in this study), *An. trinkae* and *An. oswaldoi* s.l., is of significance to vector control personnel in Colombia, as it could result in wasted resources, with unnecessary control measures mistakenly being applied against this non-vector species.

Topotypic ITS2 sequences of *An. dunhami* were less than 1% dissimilar to those reported by Calado et al. (2008). The two remaining GenBank entries labelled as *An. dunhami* are problematic: U9236 (Tefé) (Lounibos et al. 1998) and AF462378 (Acre, Brazil) (Marrelli et al. 2005). As highlighted by Calado et al. (2008), signifi-

cant differences exist between the GenBank submission (U9236) and the published sequences in the paper of Lounibos et al. (1998) and the submitted sequence shows 15 base differences (all indels) from our *An. dunhami* sequences. An author of the present study (JEC) and co-author of Lounibos et al. (1998) confirms that manual sequencing errors are most likely to account for these discrepancies. GenBank entry AF462378 of Marrelli et al. (2005) was more similar (96%) to *An. oswaldoi* s.s. from Brazil, São Paulo (AF425915), than topotypic *An. dunhami* 86% (68/496 variant bases).

Although the two *An. dunhami* specimens detected were field identified by females as *An. nuneztovari* s.l., these taxa are easily differentiated using either ITS2 or COI sequence data or male genitalia (Calado et al. 2008). *An. dunhami* and *An. nuneztovari* C share only 96.9% identity at ITS2 (Fig. 1) and 97.5% at COI (Fig. 2). *An. dunhami* ITS2 sequences generated in this research share 78–80% identity with *An. trinkae* sequences in GenBank (U92346, U92355 and Y09075) (Fritz 1998, Lounibos et al. 1998).

Contrary to the low level of intra-specific COI sequence divergence, the mean inter-specific K2P distance between *An. dunhami*, *An. nuneztovari* and *An. trinkae* was 0.067 (range 0.025–0.093). Comparable results were reported in other barcode studies on mosquitoes, i.e. 0.058 between *Anopheles fluviatilis* and *Anopheles minimus* (Kumar et al. 2007) and 10.4% (range 0.2–17.2%) for nine genera of Culicidae (Cywinska et al. 2006). In contrast to the ITS2 dataset, using mtDNA markers, *An. dunhami* and *An. nuneztovari* appear most closely related with 2.5% sequence divergence, but significantly higher than intra-specific comparisons.

Herein we verified the presence of *An. dunhami* in Colombia for the first time using both ITS2 sequences and COI barcodes. Although *An. dunhami* has not been implicated in malaria transmission (Peyton 1993, Lounibos et al. 1998), adult females, the stage most readily collected in epidemiological studies, can be easily misidentified as *An. nuneztovari* s.l., *An. oswaldoi* s.l. or *An. trinkae*, all of which are proven natural vectors of malaria in South America (Faran 1980, Hayes et al. 1987, Herrera et al. 1987, Olano et al. 2001, Quiñones et al. 2006, Gutiérrez et al. 2009). It is possible that *An. dunhami* may also be a competent vector; however, its role in local malaria transmission is masked by the presence of other morphologically similar species, which are traditionally known as malaria vectors. Knowledge of mosquito species present in vector control regions is important. In Colombia, local vector control strategies vary according to the vector species present and this in turn is driven by the entomological surveys, thus it is now imperative that morphological keys in Colombia are updated to include *An. dunhami* and to determine novel morphological characters to differentiate this species from *An. nuneztovari* in the adult female stage if possible. It seems likely that this species will also be present in the Amazonian Region of Peru and that its eventual known distribution could include a large portion of the Amazon Basin. Topotypic ITS2 and COI sequences of *An. dunhami* from Tefé generated in this study will facilitate the identification of this species across its geographical range.




Fig. 2: bootstrap neighbor-joining tree constructed using 1,000 replicates of Kimura two-parameter distance model genetic distance matrices of cytochrome c oxidase unique haplotypes (680 bp) belonging to 23 specimens (only unique haplotypes are shown) of *Anopheles nuneztovari* [NortedeSantander, Colombia (n=10), GenBank HQ315868–HQ315877], *Anopheles dunhami* [Kilómetro 12, Amazonas, Colombia (n = 1), HQ315867; Tefé, Amazonas, Brazil (n = 4), HQ315878–HQ315881; EU848330; EU848330 (same haplotype as EU848329), EU848331, EU848332 (Calado et al. (2008))]. Out-groups: *Anopheles trinkae* (Sardina Yacu, Napo State, Ecuador, HQ315882–HQ315883), *Anopheles oswaldoi* B (HQ315866) and *Anopheles rangeli* (HQ315865).

ACKNOWLEDGEMENTS

The authors are especially grateful to La Secretaria de Salud del Amazonas, Colombia. Parts of this research were performed at the Smithsonian Institution under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. The material contained within this manuscript has been reviewed by the Walter Reed Army Institute of Research. There is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the authors, and are not to be construed as official, or as reflecting true views of the Department of the Army or the Department of Defense.

REFERENCES

- Calado DC, Foster PG, Bergo ES, dos Santos CL, Galardo AK, Sallum MA 2008. Resurrection of *Anopheles goeldii* from synonymy with *Anopheles nuneztovari* (Diptera: Culicidae) and a new record for *Anopheles dunhami* in the Brazilian Amazon. *Mem Inst Oswaldo Cruz* 103: 791-799.
- Causey OR 1945. Description of *Anopheles* (*Nyssorhynchus*) *dunhami*, a new species from the upper Amazon Basin. *J Natl Malar Soc* 4: 231-234.
- Collins FH, Paskewitz SM 1996. A review of the use of ribosomal DNA (rDNA) to differentiate among cryptic *Anopheles* species. *Insect Mol Biol* 5: 1-9.
- Cywinska A, Hunter FF, Hebert PD 2006. Identifying Canadian mosquito to species through DNA barcodes. *Med Vet Entomol* 20: 413-424.
- Faran ME 1980. Mosquito studies (Diptera: Culicidae) XXXIV. A revision of the Albimanus section of the subgenus *Nyssorhynchus* of *Anopheles*. *Cont Am Entomol Inst* 15: 1-215.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3: 294-299.
- Fritz GN 1998. Sequence analysis of the rDNA internal transcribed spacer 2 of five species of South American human malaria mosquitoes. *DNA Seq* 8: 215-221.
- Fritz GN, Conn J, Cockburn A, Seawright J 1994. Sequence analysis of the ribosomal DNA internal transcribed spacer 2 from populations of *Anopheles nuneztovari* (Diptera: Culicidae). *Mol Biol Evol* 11: 406-416.
- González R, Carrejo NS 2007. *Introducción al estudio taxonómico de Anopheles de Colombia. Claves y notas de distribución*, Universidad del Valle, Cali, 237 pp.
- Gutiérrez LA, González JJ, Gómez GF, Castro MI, Rosero DA, Luckhart S, Conn JE, Correa MM 2009. Species composition and natural infectivity of anthropophilic *Anopheles* (Diptera: Culicidae) in the states of Córdoba and Antioquia, Northwestern Colombia. *Mem Inst Oswaldo Cruz* 104: 1117-1124.
- Hayes J, Calderon G, Falcon R, Zambrano V 1987. Newly incriminated anopheline vectors of human malaria parasites in Junin Department, Peru. *J Am Mosq Control Assoc* 3: 418-422.
- Hebert PD, Cywinska A, Ball SL, deWaard JR 2003. Biological identifications through DNA barcodes. *Proc Biol Sci* 270: 313-321.
- Herrera S, Suárez MF, Sanchez GI, Quiñones ML, de Herrera M 1987. Uso de la técnica radioinmunoensayo IRMA en *Anopheles* de Colombia para la detección de esporozoitos de *Plasmodium*. *Colomb Med* 18: 2-6.
- Kimura M 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16: 111-120.
- Kumar NP, Rajavel AR, Natarajan R, Jambulingam P 2007. DNA barcodes can distinguish species of Indian mosquitoes (Diptera: Culicidae). *J Med Entomol* 44: 1-7.
- Kumar S, Nei M, Dudley J, Tamura K 2008. MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Brief Bioinform* 9: 299-306.
- Lane J 1953. *Neotropical Culicidae*, Vol. 1, University of São Paulo, São Paulo, 548 pp.
- Linton Y-M, Harbach RE, Anthony TG, Chang MS, Asmad M 2001. Morphological and molecular identity of *Anopheles* (*Cellia*) *sundaicus* (Diptera: Culicidae), the nominotypical member of a malaria vector species complex in Southeast Asia. *Syst Entomol* 26: 357-366.
- Lounibos LP, Wilkerson RC, Conn JE, Hribar LJ, Fritz GN, Danoff-Burg JA 1998. Morphological, molecular and chromosomal discrimination of cryptic *Anopheles* (*Nyssorhynchus*) (Diptera: Culicidae) from South America. *J Med Entomol* 35: 830-838.
- Maddison WP, Maddison DR 2003. *MacClade: analysis of phylogeny and character evolution, version 4.06*, Sinauer Associates, Sunderland, MA.
- Marrelli MT, Floeter-Winter LM, Malafronte RS, Tadei WP, Lourenço-de-Oliveira R, Flores-Mendoza C, Marinotti O 2005. Amazonian malaria vector anopheline relationships interpreted from ITS2 rDNA sequences. *Med Vet Entomol* 19: 208-218.
- Olano VA, Brochero HL, Sáenz R, Quiñones ML, Molina JA 2001. Mapas preliminares de la distribución de especies de *Anopheles* vectores de malaria en Colombia. *Biomedica (Bogotá)* 21: 402-408.
- Peyton EL 1993. *Anopheles* (*Nyssorhynchus*) *dunhami*, resurrected from synonymy with *Anopheles nuneztovari* and validated as a senior synonym of *Anopheles trinkae* (Diptera: Culicidae). *Mosq Syst* 25: 151-156.
- Quiñones ML, Ruiz F, Calle DA, Harbach RE, Erazo HF, Linton Y-M 2006. Incrimination of *Anopheles* (*Nyssorhynchus*) *rangeli* and *An. (Nys.) oswaldoi* as natural vectors of *Plasmodium vivax* in Southern Colombia. *Mem Inst Oswaldo Cruz* 101: 617-623.
- Ruiz F 2010. *Systematics of the Anopheles* (*Nyssorhynchus*) *oswaldoi* group in Latin America, PhD Thesis, Canterbury Christ Church University, Canterbury, 189 pp.
- Saitou N, Nei M 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406-425.
- Sierra DM, Velez ID, Linton Y-M 2004. Malaria vector *Anopheles* (*Nyssorhynchus*) *nuneztovari* comprises one genetic species in Colombia based on homogeneity of nuclear ITS2 rDNA. *J Med Entomol* 41: 302-307.
- Suárez MF, Quiñones M, Fleming GA, Robayo M 1988. *Guía introductoria a la morfología de Anopheles y clave para determinación de las principales especies de Colombia*, DCD, Ministerio de Salud, Bogotá, 120 pp.
- Swofford DL 2003. *PAUP*: phylogenetic analysis using parsimony, version 4.0b 10*, Sinauer Associates, Sunderland, MA.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG 1997. The Clustal-X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25: 4876-4882.